


AD-A144 760

TWO SILICON PHTHALOCYANINES AND A SILICON  
NAPHTHALOCYANINE SYNTHESIS ELEC..(U) CASE WESTERN  
RESERVE UNIV CLEVELAND OHIO DEPT OF CHEMISTRY  
B L WHEELER ET AL. 15 AUG 84 CWRU/DC/TR-14 F/G 7/4

1/1

UNCLASSIFIED

NL




END

DATE

FILMED

10-84

DTIC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A144 760

DTIC FILE COPY

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

12

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report CWRU/DC/TR14	2. GOVT ACCESSION NO. <b>AD-A144 760</b>	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Two Silicon Phthalocyanines and a Silicon Naphthalocyanine: Synthesis, Electrochemistry and Electrogenerated Chemiluminescence		5. TYPE OF REPORT & PERIOD COVERED
7. AUTHOR(s) Bob L. Wheeler, G. Nagasubramanian, Allen J. Bard, Lee A. Schechtman, David R. Dininny and Malcolm E. Kenney		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry Case Western Reserve University Cleveland, OH 44106		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0693
11. CONTROLLING OFFICE NAME AND ADDRESS Department of the Navy Office of Naval Research Arlington, VA 22217		10. PROGRAM ELEMENT PROJECT TASK AREA & WORK UNIT NUMBERS NR 356-590
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE August 15, 1984
		13. NUMBER OF PAGES 35
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)  <div style="text-align: right;"> <b>DTIC</b>  <b>ELECTE</b>  <b>AUG 24 1984</b> </div>		
18. SUPPLEMENTARY NOTES  Accepted for publication in J. Am. Chem. Soc.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Silicon Phthalocyanines, Silicon Naphthalocyanines, Oligomeric Silicon Phthalocyanines, Electrochemistry, Electrogenerated Chemiluminescence		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  The synthesis, spectral characterization, and electrochemical behavior of bis- (tri-n-hexylsiloxy)-2,3-phthalocyaninosilicon $[\text{SiPc}(\text{OR})_2]$ , its dimer $[\text{RO}(\text{SiPcO})_2\text{R}]$ , and its naphthalocyanine analog $[\text{SiNc}(\text{OR})_2]$ are described. All compounds show near UV absorption corresponding to Soret and N bands and in- tense absorption in the visible-near IR region corresponding to Q-bands. In $\text{CH}_2\text{Cl}_2$ , within the solvent stability limit, there are two reductions and one oxidation for $\text{SiPc}(\text{OR})_2$ and two reductions and two oxidations for $\text{RO}(\text{SiPcO})_2\text{R}$ and $\text{SiNc}(\text{OR})_2$ ; all appear as reversible one electron waves, although $n=2$ for		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

the dimer. The difference in the peak potentials of the first oxidation and first reduction waves agrees well with the excitation energy and fluorescence (corresponding to Q-bands) of  $\text{SiPc(OR)}_2$  and  $\text{SiNc(OR)}_2$ . Both compounds emit upon electrochemical generation of reduced and oxidized forms (ecl). The corrected ecl maxima are near 684 nm for  $\text{SiPc(OR)}_2$  and 792 nm for  $\text{SiNc(OR)}_2$ ; the latter corresponds to the longest wavelength ecl emission reported so far. The phthalocyanine dimer,  $\text{RO(SiPcO)}_2\text{R}$ , is easier to reduce and easier to oxidize than the monomer. The voltammetric waves in this case involve two-electron transfers with peak splittings characteristic of 1e waves suggesting that the two phthalocyanine rings in the dimer do not strongly interact. No fluorescence or ecl was observed from the dimer.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

OFFICE OF NAVAL RESEARCH

Contract N00014-75-C-0693

Task No. NR 356-590

TECHNICAL REPORT NO. 14

Two Silicon Phthalocyanines and a Silicon Naphthalocyanine:  
Synthesis, Electrochemistry and Electrogenerated Chemiluminescence

by

Bob L. Wheeler, G. Nagasubramanian, Allen J. Bard,  
Lee A. Schechtman, David R. Dininny and Malcolm E. Kenney

Accepted

by the

Journal of the American Chemical Society

Chemistry Department  
Case Western Reserve University  
Cleveland, OH 44106

August 15, 1984

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

This document has been approved for public release  
and sale; its distribution is unlimited.

## (Abstract)

The synthesis, spectral characterization, and electrochemical behavior of bis(tri-n-hexylsiloxy)-2,3-phthalocyaninatosilicon  $[\text{SiPc}(\text{OR})_2]$ , its dimer  $[\text{RO}(\text{SiPcO})_2\text{R}]$ , and its naphthalocyanine analog  $[\text{SiNc}(\text{OR})_2]$  are described. All compounds show near UV absorption corresponding to Soret and N bands and intense absorption in the visible-near IR region corresponding to Q-bands. In  $\text{CH}_2\text{Cl}_2$ , within the solvent stability limit, there are two reductions and one oxidation for  $\text{SiPc}(\text{OR})_2$  and two reductions and two oxidations for  $\text{RO}(\text{SiPcO})_2\text{R}$  and  $\text{SiNc}(\text{OR})_2$ ; all appear as reversible one electron waves, although  $n=2$  for the dimer. The difference in the peak potentials of the first oxidation and first reduction waves agrees well with the excitation energy and fluorescence (corresponding to Q-bands) of  $\text{SiPc}(\text{OR})_2$  and  $\text{SiNc}(\text{OR})_2$ . Both compounds emit upon electrochemical generation of reduced and oxidized forms (ecl). The corrected ecl maxima are near 684 nm for  $\text{SiPc}(\text{OR})_2$  and 792 nm for  $\text{SiNc}(\text{OR})_2$ ; the latter corresponds to the longest wavelength ecl emission reported so far. The phthalocyanine dimer,  $\text{RO}(\text{SiPcO})_2\text{R}$ , is easier to reduce and easier to oxidize than the monomer. The voltammetric waves in this case involve two-electron transfers with peak splittings characteristic of 1e waves suggesting that the two phthalocyanine rings in the dimer do not strongly interact. No fluorescence or ecl was observed from the dimer.

(end of abstract)



Information For	
DATE	<input checked="" type="checkbox"/>
TIME	<input type="checkbox"/>
APPROVED	<input type="checkbox"/>
BY	
Classification/	
Availability Codes	
and/or	
Dist	
<b>A-1</b>	

## INTRODUCTION

Phthalocyanine compounds often show high thermal and chemical stability and interesting optical and electrical properties.<sup>1-5</sup> We and others have been interested in the conductivities of these materials<sup>6,7</sup> and in their application to the sensitization and stabilization of semiconductor electrodes in photoelectrochemical cells.<sup>8,9</sup> The group IV metal phthalocyanines are of interest because various groups can be attached to the axial (or trans) positions.<sup>10</sup> Further, some group IV phthalocyanines are linear stacked polymers.<sup>4,11</sup> We describe here the synthesis and the electrochemical and spectroscopic properties of a trialkylsiloxysilicon phthalocyanine, bis(tri-*n*-hexylsiloxy)phthalocyaninosilicon  $[\text{SiPc}(\text{OR})_2]$ , and of its 2,3-naphthalocyanine analog,  $[\text{SiNc}(\text{OR})_2]$ , (Fig. 1). We also describe the synthesis and properties of the dimeric analog of the phthalocyanine  $[\text{RO}(\text{SiPcO})_2\text{R}]$ . The presence of the trialkylsiloxy groups on the central Si atom leads to relatively high solubility in these compounds and permits studies of solutions of them at the mM level.

## EXPERIMENTAL

### Synthesis of $\text{SiNc}[\text{OSi}(\text{n-C}_6\text{H}_{13})_3]_2$ and $\text{SiPc}[\text{OSi}(\text{n-C}_6\text{H}_{13})_3]_2$

2,3-Naphthalenedicarbonitrile. The procedure employed is a modification of one used by Luk'yanets.<sup>12</sup> A mixture of  $\alpha, \alpha, \alpha', \alpha'$ -tetrabromo-*o*-xylene (200 g), fumaronitrile (64 g), NaI (200 g), and dry dimethylformamide (2L) was heated at 75° for 5.5 h. The resultant was allowed to cool to 65°C and then poured into a mixture of ice and water (3L). Sodium bisulfite was added to the suspension obtained (which was dark red), until it turned yellow-tan (65 g). The product was filtered off, washed, vacuum-dried (70° C), and weighed (65

g, 78%).

A small portion of this product was sublimed (160° C) in air at atmospheric pressure: mp 256° C (lit<sup>12</sup> 251° C); IR (Nujol) 2220 (C≡N) cm<sup>-1</sup>; NMR (200 MHz, CDCl<sub>3</sub>) δ 8.37 (s, 1,4-H); 8.00 (m, 5,8-H), 7.81 (m, 6,7-H); mass spectrum, m/z (rel intensity) 178 (M<sup>+</sup>, 100), 151 (20). Anal. Calcd. for C<sub>12</sub>H<sub>6</sub>N<sub>2</sub>: C, 80.88; H, 3.40; N, 15.72. Found: C, 80.83; H, 3.56; N, 15.68.

The sublimed compound is composed of small, colorless needles. The synthesis product is a beige color and is satisfactory for use as an intermediate.

1,3-Diiminobenz(f)isoindoline. The synthesis used by Esposito for 1,3-diiminoisoindoline<sup>13</sup> served as a model. Anhydrous NH<sub>3</sub> was slowly bubbled through a stirred mixture of 2,3-naphthalenedicarbonitrile (10.2 g), sodium methoxide (1.5 g), and dry methanol (90 mL) for 45 min. With continued NH<sub>3</sub> introduction, the mixture was refluxed for 3 h. After the resultant had cooled, the product was collected by filtration, washed, air-dried, and weighed (9.2 g, 82%): IR (Nujol) 3285, 3200, 3080, 1700, 1640, 1545, 1520 cm<sup>-1</sup>; NMR (100 MHz, (CD<sub>3</sub>)<sub>2</sub>SO), δ 8.31 (s, 1,4-H), 8.08 (m, 5,8-H), 7.68 (m, 6,7-H); mass spectrum m/z (rel intensity) 196 (70), 195 (M<sup>+</sup>, 100), 179 (68), 126 (51), 63 (59), 51 (55).

From batch to batch the product ranges from yellow to green. Attempts made to purify it by recrystallization and sublimation failed. As obtained, the product is sufficiently pure for use as a naphthalocyanine precursor.

SiNcCl<sub>2</sub>. A mixture of 1,3-diiminobenz(f)isoindoline (6.0 g), SiCl<sub>4</sub> (5.4 mL), dry tetrahydronaphthalene (40 mL), and dry tri-n-butylamine (20 mL) was refluxed for 2.5 h. The resultant was allowed to cool and then diluted with methanol (30 mL). The product was filtered off, washed, air-dried, and weighed (5.8 g): IR (Nujol) 475 (m, Si-Cl) cm<sup>-1</sup>.



The product is quite impure, containing perhaps as much as 20-40% impurities. It is dark green and has very low solubility in a variety of organic solvents. It also has little volatility. Attempts made to purify it failed.

SiNc(OH)<sub>2</sub>. A mixture of SiNcCl<sub>2</sub> (5.8 g) and concentrated H<sub>2</sub>SO<sub>4</sub> (200 mL) was stirred for 2 h, and then poured over ice (600 g). The solid was filtered off, washed with water and an acetone-water solution. A mixture of this solid and concentrated NH<sub>4</sub>OH (150 mL) was refluxed for 1 h and allowed to cool. The resulting solid was filtered off, washed, vacuum-dried, and weighed (5.6 g). Part of it (4.3 g) was extracted with pyridine in a Soxhlet extractor, washed, vacuum-dried, and weighed (2.8 g).

A pure sample of the hydroxide was obtained by hydrolysis of SiNc[OSi(n-C<sub>6</sub>H<sub>13</sub>)<sub>3</sub>]<sub>2</sub> (see below). A mixture of chromatographed SiNc[OSi(n-C<sub>6</sub>H<sub>13</sub>)<sub>3</sub>]<sub>2</sub> (100 mg), water (3 mL), 1,2-dimethoxyethane (60 mL), and concentrated HCl (0.06 mL) was refluxed for 5 h. The resulting suspension was allowed to cool, and the solid was collected by filtration, washed, vacuum-dried, and weighed (29 mg, 50%): IR(Nujol) 3435 (O-H), 840 (Si-O) cm<sup>-1</sup>. Anal. Calcd. for C<sub>48</sub>H<sub>26</sub>N<sub>8</sub>O<sub>2</sub>Si: C, 74.40; H, 3.38; Si, 3.62. Found: C, 74.04; H, 3.71; Si, 3.52.

The compound is green. It is, at best, only slightly soluble in a variety of organic solvents. The product of the SiNcCl<sub>2</sub> synthesis is generally satisfactory for use as an intermediate.

SiNc[OSi(n-C<sub>6</sub>H<sub>13</sub>)<sub>3</sub>]<sub>2</sub>. For the synthesis of this compound, a mixture of SiNc(OH)<sub>2</sub> made from SiNcCl<sub>2</sub> (600 mg), tri-n-hexylchlorosilane (3.0 mL), dry tri-n-butylamine (2.0 mL), and dry 3-picoline (70 mL) was refluxed for 1.5 h. The solution obtained was allowed to cool and then filtered (no residue). The filtrate was poured into an ethanol-water solution (1:1, 100 mL), and the product was separated by filtration from the resulting

suspension. After being washed, it was vacuum-dried, and weighed (808 mg, 78%).

The product was chromatographed on alumina (activity I) with a toluene-hexanes solution (3:1) as the eluant: mp 278° C; IR (Nujol) 1250 (Si-CH<sub>2</sub>), 1040 (Si-O-Si) cm<sup>-1</sup>; NMR (200 MHz, CDCl<sub>3</sub>) δ 10.11 (s, 1,4-Nc), 8.69 (m, 5,8-Nc), 7.94 (m, 6,7-Nc), 0.61 (m, ε-CH<sub>2</sub>), 0.41 (t, CH<sub>3</sub>), 0.22 (m, δ-CH<sub>2</sub>), 0.06 (m, γ-CH<sub>2</sub>), -0.98 (m, β-CH<sub>2</sub>), -2.05 (m, α-CH<sub>2</sub>). Anal. Calcd. for C<sub>84</sub>H<sub>102</sub>N<sub>8</sub>O<sub>2</sub>Si<sub>3</sub>: C, 75.29; H, 7.67; Si, 6.29. Found: C, 75.05; H, 7.69; Si, 6.03.

The compound is green. It is stable and soluble in a variety of nonpolar and chlorinated organic solvents. However, it is insoluble in acetonitrile. As shown by its NMR spectrum, the synthesis product contains a small amount of the dimer (n-C<sub>6</sub>H<sub>13</sub>)<sub>3</sub>SiO(SiNcO)<sub>2</sub>Si(n-C<sub>6</sub>H<sub>13</sub>)<sub>3</sub>.

SiPc[OSi(n-C<sub>6</sub>H<sub>13</sub>)<sub>3</sub>]<sub>2</sub>. A synthetic procedure similar to that described by Douglass<sup>14</sup> was followed. A mixture of SiPc(OH)<sub>2</sub> (0.75 g), tri-n-hexylchlorosilane (5.0 mL), and dry pyridine (75 mL) was refluxed for 5 h. After the solution obtained had been allowed to cool, it was filtered (no residue) and concentrated under vacuum. The resulting oil was mixed with pentanes (10 mL) and the slurry formed was filtered. The solid was washed with pentanes (10 mL) and an acetone-water solution, vacuum-dried, and weighed (0.78 g). To recover additional product, the pentane filtrates were combined, concentrated, and filtered. The solid was washed, vacuum-dried, and weighed (0.28 g, combined yield 78%).

The combined solids were chromatographed as described by Janson<sup>15</sup> on alumina (activity I) with a toluene-hexanes solution (1:2) as the eluant: mp 175° C (lit<sup>15</sup> 175° C); IR (Nujol) 1250 (Si-CH<sub>2</sub>), 1040 (Si-O-Si) cm<sup>-1</sup>; NMR (200 MHz, CDCl<sub>3</sub>) δ 9.63 (m, 3,6-Pc), 8.31 (m, 4,5-Pc), 0.82 (m, ε-CH<sub>2</sub>), 0.71 (t, CH<sub>3</sub>), 0.36 (m, δ-CH<sub>2</sub>), 0.02 (m, γ-CH<sub>2</sub>), -1.28 (m, β-CH<sub>2</sub>), -2.45 (m, α-CH<sub>2</sub>). Anal. Calcd. for C<sub>68</sub>H<sub>94</sub>N<sub>8</sub>O<sub>2</sub>Si<sub>3</sub>: C, 71.66; H, 8.31; Si, 7.39.

Found: C, 71.60; H, 8.37; Si, 7.41.

In accordance with expectations, the compound is stable and is blue when finely divided. It can be sublimed under vacuum at elevated temperatures and is soluble in many organic solvents including mixed pentanes. The synthesis product contains, as is evident from its NMR spectrum, a small amount of  $(n\text{-C}_6\text{H}_{13})_3\text{SiO}(\text{SiPcO})_2\text{Si}(n\text{-C}_6\text{H}_{13})_3$ .<sup>14</sup>

$(n\text{-C}_6\text{H}_{13})_3\text{SiO}(\text{SiPcO})_2\text{Si}(n\text{-C}_6\text{H}_{13})_3$ .<sup>14</sup> The procedure used earlier<sup>16</sup> for a similar dimer was employed as a model. A mixture of  $\text{SiPc}(\text{OH})_2$  (540 mg),  $\text{SiPcCl}_2$  (294 mg), dry tri-*n*-butylamine (1.0 mL), and dry, nitrogen-purged quinoline (65 mL) was refluxed for 15 min. The resulting suspension was allowed to cool, mixed with concentrated HCl (46 mL), and filtered. The solid obtained was washed. A mixture of this solid and concentrated  $\text{H}_2\text{SO}_4$  (50 mL) was stirred for 2 h, and then poured over ice (200 g). The solid was filtered off and washed. A mixture of the solid, concentrated  $\text{NH}_4\text{OH}$  (150 mL), and pyridine (10 mL) was refluxed for 1 h, cooled, and filtered. The resulting hydroxy-capped oligomers were washed, vacuum-dried, and weighed (763 mg, 96%).

A mixture of the oligomers (707 mg), tri-*n*-hexylchlorosilane (0.5 mL), and dry pyridine (125 mL) was refluxed for 2.5 h. The solution obtained was filtered while hot (46 mg residue) and evaporated to dryness under vacuum. After being washed, the solid was air-dried and weighed (1.13 g).

This solid was chromatographed on alumina (activity III). The desired product was eluted with a toluene-hexanes solution (1:3) (376 mg, 32% overall): mp 201° C; IR(Nujol) 1250 (Si-CH<sub>2</sub>), 1025 (SiR-O-SiPc), 970 (SiPc-O-SiPc) cm<sup>-1</sup>; NMR (200 MHz, CDCl<sub>3</sub>)  $\delta$  9.00 (m, 3,6-Pc), 8.31 (m, 4,5-Pc), 0.43 (overlapping m, CH<sub>3</sub> and  $\epsilon$ -CH<sub>2</sub>), -0.20 (m,  $\delta$ -CH<sub>2</sub>), -0.65 (m,  $\gamma$ -CH<sub>2</sub>), -2.29 (m,  $\beta$ -CH<sub>2</sub>), -3.54 (m,  $\alpha$ -CH<sub>2</sub>).

The compound is a blue solid that is soluble in common organic

solvents. Further elution of the column yielded the trimer, the tetramer, and a mixture with a substantial pentamer content.

Elemental analyses were done by Galbraith Laboratories, Inc., Knoxville, TN.

#### Electrochemical Measurements

The solvent, methylene chloride ( $\text{CH}_2\text{Cl}_2$ , MCB Omnisolv grade; EM Science, Gibbstown, NJ), was dried by vacuum distillation over molecular sieves and stored in a helium-filled Vacuum Atmosphere Corporation (Hawthorne, CA) glove box. Polarographic grade tetra-n-butylammonium perchlorate (TBAP), which was recrystallized twice from acetone-ether and dried under a vacuum of better than  $10^{-5}$  torr for two days, was used as the supporting electrolyte at 0.1 M concentration. For electrochemical measurements a three compartment electrochemical cell of 25 mL capacity was used. A Pt disc of  $0.03 \text{ cm}^2$  area, was used as the working electrode. A large area ( $>40 \text{ cm}^2$ ) Pt gauze, separated from the working electrode by a fine porosity glass frit, was the counter electrode. The reference electrode, a quasireference Ag electrode, denoted as AgRE, was also separated from the main compartment by a fine porosity glass frit and the reference and counter electrodes were immersed in a solution containing the solvent,  $\text{CH}_2\text{Cl}_2$ , and 0.1 M TBAP. In a typical experiment, before the addition of the phthalocyanine or naphthalocyanine compounds, the  $\text{CH}_2\text{Cl}_2$ , 0.1 M TBAP solution was stirred with activated alumina (Woelm Neutral Alumina - activity Super I, Woelm Pharma GmbH and Co., Eschwege, BRD) to remove any residual water, and was then placed in all three cell compartments. The concentrations of the  $\text{SiPc}(\text{OR})_2$ ,  $\text{RO}(\text{SiPcO})_2\text{R}$ , and  $\text{SiNc}(\text{OR})_2$  in the working electrode compartment were around 2.2 mM, 1.0 mM, and 1.4 mM, respectively. All the experiments were carried out inside the glove box. At the conclusion of the experiments the cell was taken out of the glove box, and the potential of the AgRE was measured

against the aqueous saturated calomel electrode (SCE). Because of the slow shift in the AgRE, mainly due to a seepage of redox couple into the reference compartment, potentials reported vs. SCE were within  $\pm 100$  mV for  $\text{SiNc(OR)}_2$  and  $\pm 10$  mV for  $\text{SiPc(OR)}_2$  and  $\text{RO(SiPcO)}_2\text{R}$ . Positive feedback iR compensation was used to minimize the uncompensated resistance during cyclic voltammetric experiments.

A Princeton Applied Research (PAR) model 173 potentiostat and a PAR model 175 universal programmer equipped with a Houston Instruments (Austin, TX) model 2000 X-Y recorder was used to obtain cyclic voltammograms and, with a time base module, for chronoamperometry. For ac voltammetry, a Soltec (Sun Valley, CA) model 6432, X-Y<sub>1</sub>Y<sub>2</sub> recorder was employed. For the a-c voltammetry, the lock-in amplifier technique which yields the in-phase ( $0^\circ$  component) and the out-of-phase ( $90^\circ$  component) components of a sine wave superimposed on a linear potential ramp was employed. For the above measurement a PAR model 5204 lock-in amplifier was used, which output simultaneously the phase angle and the in-phase component. The low harmonic distortion ac signal ( $12 \text{ mV}_{\text{pk-pk}}$ ) at different frequencies was obtained for input into the potentiostat from a model 200 CD wide range oscillator (Hewlett-Packard, Palo Alto, CA).

#### Spectral Studies

The electronic spectra of the two compounds were recorded with a Cary 17D (Varian Associates, Sunnyvale, CA) spectrophotometer. Infrared spectra were obtained with a Perkin-Elmer 598 infrared spectrophotometer (Perkin-Elmer Corp., Norwalk, CT). A Varian HA-100 NMR spectrometer equipped for Fourier-transform operation was used to record the 100 MHz spectrum while a Varian XL-200 instrument was used to record the 200 MHz

spectra. Mass spectra were obtained with an AEI MS-30 spectrometer (Associated Electrical Industries, Manchester, England). The samples were introduced by means of a heated solid probe (350° C) and ionized by electron impact (70 eV). For fluorescence studies an Aminco-Bowman spectrophotometer with appropriate slits to achieve a band pass of 5 nm for both excitation and emission was used. A photomultiplier tube R928 (Hamamatsu TV Co., Ltd., Middlesex, NJ) in conjunction with a grating monochromator model 7240 (Oriel Corporation, Stamford, CT) with 1  $\mu$ m blaze and slits for 30 nm bandwidth were used for recording the ecl spectra. The procedure used was similar to that reported previously.<sup>17</sup>

## RESULTS AND DISCUSSION

### Synthetic Considerations

The syntheses for  $\text{SiNcCl}_2$  and  $\text{SiNc(OR)}_2$  and the synthesis for  $\text{SiNc(OH)}_2$  from  $\text{SiNcCl}_2$  are modeled on analogous syntheses for the corresponding phthalocyanines.<sup>13,14,18</sup> In view of the nature of the routes previously found to be useful for the preparation of silicon phthalocyanines,<sup>19</sup> and the satisfactory character of the synthesis for  $\text{SiNcCl}_2$ , it seems probable that  $\text{SiNcCl}_2$  will generally be found to be a good entry point for the synthesis of silicon naphthalocyanines. Further, since the synthesis for  $\text{SiNc(OH)}_2$  is also satisfactory, it appears the  $\text{SiNcCl}_2$  and  $\text{SiNc(OH)}_2$  will be generally found to be good intermediates for the preparation of silicon naphthalocyanines.

### Chemical and Physical Properties.

The chemical stability of the SiNc system is illustrated by its ability to survive intact during the concentrated  $\text{H}_2\text{SO}_4$  hydrolysis of  $\text{SiNcCl}_2$ . In large part the resistance of the ring in this system to degradation can be attributed to its aromaticity while the resistance of the silicon to expulsion from the system can be associated with the macrocyclic (chelate)

effect. The nonlability of the silicon of this system is of interest because the silicon is bound by Si-N bonds. The parallel nonlability of the silicon in the SiPc system has previously been noted as being of interest for the same reason.<sup>18</sup>

On the basis of the available data, it is clear that the solubility, volatility, and melting point characteristics of  $\text{SiNcCl}_2$ ,  $\text{SiNc(OH)}_2$ , and  $\text{SiNc[OSi(n-C}_6\text{H}_{13})_3]_2$  are directly associated with the size and nature of their axial ligands. These properties can be understood in terms of the relative strengths of the  $\pi$ - $\pi$  and other intermolecular interactions involved.

In accordance with expectations, the shapes and positions of the naphthalocyanine bands in the infrared spectra of the three naphthalocyanines are quite similar. Also in accordance with expectations, the pattern of the naphthalocyanine bands is, as is illustrated by the spectrum of  $\text{SiNc(OH)}_2$ , Fig. 2, comparatively simple.

The NMR spectrum of  $\text{SiNc[OSi(n-C}_6\text{H}_{13})_3]_2$ , Fig. 3, shows that the naphthalocyanine ring produces, not surprisingly, a large ring-current effect. Because of this, the assignments of the individual resonances of the hexyl compound are straightforward. However, it is of interest that although the hexyl resonances of this compound and those of  $\text{SiPc[OSi(n-C}_6\text{H}_{13})_3]_2$  are similarly positioned, the resonances of the inner hexyl protons of the naphthalocyanine are shifted upfield less than the resonances of the corresponding protons of the phthalocyanine. Conversely, the resonances of the outer hexyl protons of the naphthalocyanine are shifted more. Since the chains are probably similarly extended and positioned in the two compounds (in solution), it appears that for comparable regions above the rings and near their centers the ring-current effect is somewhat less for the naphthalocyanine ring, but that for

comparable regions above and further out the effect is somewhat greater. The greater ring-current effect of the naphthalocyanine ring further out can be ascribed to its greater spread.

Comparison of the positions of like protons in the hexyl chains of  $\text{SiPc}(\text{OSi}(\text{n-C}_6\text{H}_{13})_3)_2$  and  $(\text{n-C}_6\text{H}_{13})_3\text{SiO}(\text{SiPcO})_2\text{Si}(\text{n-C}_6\text{H}_{13})_3$  shows that the hexyl protons of the dimer are upfield of those of the monomer. This is attributed to the additivity of the shielding effects of the rings in the dimer.<sup>20</sup>

### Electrochemical Measurements

The stability limits of  $\text{CH}_2\text{Cl}_2$  with 0.1 M TBAP were + 1.6 V to -2.3 V vs. AgRE. In the working potential window two reductions and one oxidation of  $\text{SiPc}(\text{OR})_2$  and two reductions and two oxidations of  $\text{RO}(\text{SiPcO})_2\text{R}$  and  $\text{SiNc}(\text{OR})_2$  were observed. Typical cyclic voltammograms at 20 mV/s are shown in Fig. 4. From their general shape, all waves appear to be reversible one-electron transfers. Such multiple oxidation and reduction waves have been reported for other phthalocyanines in nonaqueous media.<sup>16,21</sup> The average values of the peak potentials obtained at several scan rates are given in Table 1. Note that while it is easier to oxidize  $\text{SiNc}(\text{OR})_2$  than  $\text{SiPc}(\text{OR})_2$ , the reduction potentials of both are nearly the same. The lack of observation of a second oxidation wave in the  $\text{SiPc}(\text{OR})_2$  species probably can be attributed to the onset of background oxidation before its appearance. A-c voltammetry experiments performed with benzonitrile as the solvent indicate a second oxidation wave near the solvent limit at +2.1 V vs. SCE, 1.1 V positive of the first oxidation wave in this solvent. The difference in potential in  $\text{CH}_2\text{Cl}_2$  between the first and second reduction waves, 0.5 V, agrees with that found by Lever, et al.,<sup>21c</sup> for  $\text{PcSi}(\text{O-t-amyl})_2$  in DMF. Further information about the electrochemical processes were obtained by investigating the dependence of peak currents



( $i_p$ ) and potentials ( $E_p$ ) on scan rate ( $v$ ), and by ac voltammetric methods. For the first reduction peaks of all these compounds  $i_{pc}/v^{1/2}$  was constant and  $i_{pc} = i_{pa}$  for  $v$  between 0.005 and 2.0 V/s. Similar results were obtained for the first oxidation waves. These results demonstrate that the products of the first oxidation and reduction are stable on the cyclic voltammetric time scale and the waves are diffusion limited. Although it is more difficult to obtain quantitative data for the later peaks, the general shape and behavior of these also suggest stability of the product of the second reduction for all compounds and the product of the second oxidation of  $RO(SiPcO)_2R$  and of  $SiNc(OR)_2$ . The waves also appear to be Nernstian ones with peak splittings ( $\Delta E_p = E_{pa} - E_{pc}$ ) of the order of 80 mV. Deviation of  $\Delta E_p$  from the rigorous Nernstian value of about 60 mV can probably be attributed to effects of uncompensated resistance in this relatively resistive solvent. Further evidence for the rapidity of the heterogeneous electron transfer reactions of all of the waves was obtained by ac voltammetric measurements (Fig. 5). The in-phase ( $0^\circ$ ) component of the ac current and the phase angle of this current with respect to the superimposed 50 Hz potential variation are essentially equal for all waves. The deviation of the phase angle from the expected Nernstian value of  $45^\circ$  again can probably be attributed to uncompensated resistance. If the phase angle is adjusted to  $90^\circ$  in the region where no faradaic current flows ( $+0.1$  -  $-0.8$  V vs AgRE), by manipulating the lock-in phase shifter, the phase angle at the peak potentials equals  $45^\circ \pm 2^\circ$ . Similarly, the second harmonic ac voltammetric peaks<sup>22</sup>, with an applied signal at 200 Hz are all essentially equal (Fig. 6), again suggesting rapid electron transfer rates. Controlled potential coulometry at the first reduction waves of  $SiPc(OR)_2$  and  $SiNc(OR)_2$  yield  $n_{app}$ -values (faradays/mole) of 1.0 and 1.1 respectively. Cyclic voltammetry of the reduced solutions, with an initial anodic scan

showed essentially the same pattern of waves as the original solution of parent compound. Re-oxidation of these solutions consumed essentially the same number of faradays as the initial reduction and regenerated the starting solution. Parallel behavior was found for coulometric oxidation at the first waves, with  $n_{app}$ -values of 1.1 (Pc) and 1.0 (Nc). These results confirm the one-electron nature of the cyclic voltammetric waves and the high stability of the one-electron reduced and oxidized products.

For the dimer  $RO(SiPcO)_2R$ , however, bulk electrolysis gave  $n_{app} = 2.0$  for oxidation at the first anodic wave and  $n_{app} = 2.2$  for reduction at the first cathodic wave. That the reduction and oxidation products were stable was clear from voltammetric scans following bulk electrolysis as well as the ability to regenerate the original parent solution electrolytically. Thus, although the voltammetric wave shapes are characteristic of one-electron reactions (e.g.,  $\Delta E_p \approx 70$  mV), these clearly represent two-electron transfers. Waves of this type have been observed before and are ascribed to molecules which have two identical non-interacting systems.<sup>23</sup> This would suggest that there is no or little interaction between the two SiPc groups in the dimeric molecule, with each accepting or losing an electron independently. However, in previous cases of noninteracting systems the potentials for electron transfer in the multi-centered species were very near those of the monomer, while in this case the potential for oxidation of the dimer is shifted by about 0.3 V towards less positive potentials (Table I), with a smaller ( $\sim 90$  mV) shift of the first reduction wave. This shift might be attributable to an inductive (electron donating) effect of the SiPc(OR) group compared to the (OR) group. Alternatively the groups may be interacting, but a rearrangement occurs following the first electron transfer which promotes the second. Because both oxidation and reduction of the dimer show the same 2e nature with peak splittings similar to the

monomer, this latter proposal seems unlikely. In a previous study of the electrochemistry of closely related SiPc-compounds,<sup>16</sup> in which the  $(n\text{-C}_6\text{H}_{13})_3$  groups attached to the Si were replaced by  $(\text{CH}_3)_2(\text{t-Bu})$ , the oxidation and reduction waves of the dimeric species were attributed to one-electron reactions based on thin layer coulometric studies of  $10^{-6}$  M solutions. The reason for this difference in behavior is not clear and bears further investigation.

Diffusion coefficients ( $D$ ) for these compounds were estimated from Cottrell plots<sup>22</sup> (current vs.  $t^{-1/2}$  for potential steps from near the rest potential (0 V vs. SCE) to the diffusion limited region of the first oxidation and reduction waves). Linear  $i$  vs.  $t^{-1/2}$  plots with zero intercepts were obtained for data between approximately 1 and 10 sec. For  $\text{SiPc}(\text{OR})_2$ , a step from 0 to +1.2 V vs. SCE or from 0 to -1.1 vs. SCE was employed; from the slope of the  $i$  vs.  $t^{-1/2}$  curve,  $D = 5.7 \pm 1.1 \times 10^{-6} \text{ cm}^2/\text{s}$ . For  $\text{SiNc}(\text{OR})_2$ , steps of 0 to +0.8 V or 0 to -1.2 V were used and yielded  $D = 5.2 \pm 0.4 \times 10^{-6} \text{ cm}^2/\text{s}$ . This slightly smaller  $D$  value is consistent with the larger ring size in the Nc species. For  $\text{RO}(\text{SiPcO})_2\text{R}$ , steps from 0 to -1.0 V or 0 to +0.9 V gave  $D = 2.2 \pm 0.3 \times 10^{-6} \text{ cm}^2/\text{s}$  taking  $n = 2$ . This comparatively low value is consistent with the bulkier nature of the dimer.

#### Absorption and Fluorescence

The UV-visible absorption spectra for  $1.1 \times 10^{-5}$  M  $\text{SiPc}(\text{OR})_2$ ,  $1.2 \times 10^{-5}$  M  $\text{RO}(\text{SiPcO})_2\text{R}$ , and  $2.5 \times 10^{-6}$  M  $\text{SiNc}(\text{OR})_2$  in  $\text{CH}_2\text{Cl}_2$  are given in Fig. 7.  $\text{SiPc}(\text{OR})_2$  and  $\text{SiNc}(\text{OR})_2$  have the features expected on the basis of previous work.<sup>24-26</sup> The spectrum of  $\text{SiPc}(\text{OR})_2$  shows the characteristic absorption for the Soret band at 350 nm, the N band at  $\sim 270$  nm, and the

series of peaks due to the Q band transitions. The absorption in the Q-band region (550-700 nm) appears to be composed of transitions from the ground state to two excited states. The absorption in the Q band region for  $\text{SiNc(OR)}_2$  is similar. The wavelength for these absorptions are listed in Table 2.

The shift in the Soret and Q band absorptions toward higher energy in the dimer  $\text{RO(SiPcO)}_2\text{R}$ , relative to  $\text{SiPc(OR)}_2$  has been explained previously in terms of exciton coupling of neutral-excitation transitions of the two rings of the dimer.<sup>27</sup> The UV-visible absorption spectrum of  $\text{RO(SiPcO)}_2\text{R}$  has all the features of the phthalocyanine dimer of these previous works.


Fluorescence is observed from  $\text{SiPc(OR)}_2$  and  $\text{SiNc(OR)}_2$  in  $\text{CH}_2\text{Cl}_2$  solution. The uncorrected emission and excitation spectra are given in Fig. 8. Both compounds show emissions in the Q band region for excitation in either Q or the Soret absorption bands. The excitation spectra of these compounds are essentially the same as the absorption spectra and show two sets of three peaks corresponding to absorption from the ground state to two different excited states. The emission spectra are essentially mirror images of the Q band absorption spectra; the emission peaks are listed in Table 2.

A solution of  $\text{RO(SiPcO)}_2\text{R}$  exhibited very weak fluorescence. However, the excitation spectrum did not match the absorption spectrum and therefore impurities were suspected as being the emitting species. Thin layer chromatography confirmed the presence of at least two different impurities. The absence of fluorescence in the pure dimer, like the absorption spectrum, can be explained in terms of the exciton model.<sup>28</sup> The lower energy exciton state, which is forbidden and therefore not observed in the absorption spectrum, is a metastable singlet state. This allows for greatly enhanced intersystem crossing into the triplet state. As a result, the singlet

emission is quenched.

### Electrogenerated Chemiluminescence

The high stability of the oxidized and reduced forms of  $\text{SiPc}(\text{OR})_2$  and  $\text{SiNc}(\text{OR})_2$  and the fact that both show strong emission upon photoexcitation suggested that these materials were good candidates for ecl studies. There has been only one previous report of ecl from a Pc system, where very weak emission was observed from metal-free Pc.<sup>29</sup> Ecl from these compounds is of interest, because there are few systems showing emission in the red and near infra-red region of the spectrum. The energetics of the annihilation process can be estimated from the redox potentials of the relevant species. Thus, the reaction enthalpies can be estimated<sup>30</sup> as about 1.8 and 1.5 eV for the  $\text{SiPc}(\text{OR})_2$  and  $\text{SiNc}(\text{OR})_2$  systems, respectively. The energetics of the lowest 0,0 bands, as estimated from the position of the highest energy emission maximum, are 1.86 and 1.61 eV. Thus, these systems may have sufficient energy to populate the emitting state directly via the electron transfer reaction:



When a Pt electrode immersed in a 1-2 mM solution of the Pc or Nc compound in  $\text{CH}_2\text{Cl}_2$  was pulsed between potentials for formation of the +1 and -1 species (for  $\text{SiPc}(\text{OR})_2$ , between -1.2 and +1.1 V and for  $\text{SiNc}(\text{OR})_2$  between -1.3 and +0.6 V) at a frequency of 0.5 Hz, weak but easily measurable emission was observed. The ecl spectra are shown in Fig. 9. For  $\text{SiNc}(\text{OR})_2$  pulsing between potentials for production of the +2 and -2 species produced the same ecl spectra; similar behavior has been seen previously, e.g., in  $\text{Ru}(\text{bpy})_3^{2+}$ <sup>31a</sup> and rubrene systems,<sup>31b</sup> and has been attributed to recombination reactions that produce the lower oxidized or reduced

species. The ecl emissions are shifted to somewhat longer wavelengths as compared to the corresponding fluorescence spectra. This shift probably can be attributed to self-absorption of the emission in ecl, since these compounds have Q bands (absorption peak, Fig. 7) located very close to the wavelength of the onset of emission in the ecl spectrum. When corrected for these absorption effects (based on the known absorber concentration, absorptivity and path length) the corrected ecl spectra are shifted to shorter wavelengths and match the fluorescence spectra. While the energetics of the electron transfer and emission processes suggest that this may be an "energy sufficient" or "S-route" system, it is marginally so. The difference in emission intensities of the anodic and cathodic pulses, clearly seen in Fig. 9, suggests some mechanistic complications. Since both the +1 and -1 forms are quite stable, the difference might be caused by at least some contribution of a "T-route" (formation of triplets in the electron transfer reaction followed by T-T annihilation), with the +1 and -1 forms quenching the triplets at different rates. T-T annihilation has been proposed in the delayed fluorescence of metal-free Pc.<sup>32</sup> Finally the ecl of the  $\text{SiNc(OR)}_2$  system represents, to our knowledge, the longest wavelength ecl emission yet reported, although other reports of ecl in the red region have appeared.<sup>29,33</sup> No ecl was observed from  $\text{RO(SiPcO)}_2\text{R}$  upon pulsing between either the first oxidation and reduction or the second oxidation and reduction waves.

#### CONCLUSIONS

In view of the structural similarities of SiPc and SiNc systems, and in view of the similarities between the physical and chemical properties of  $\text{SiNcCl}_2$ ,  $\text{SiNc(OH)}_2$ , and  $\text{SiNc(OR)}_2$  and their phthalocyanine counterparts, it is apparent that pairs of the two types of compounds will be found to be generally similar. Thus the properties of many silicon naphthalocyanines

are predictable. In  $\text{CH}_2\text{Cl}_2$ , within the solvent stability limit, there are two reductions and one oxidation for  $\text{SiPc(OR)}_2$  and two reductions and two oxidations for  $\text{RO(SiPcO)}_2\text{R}$  and  $\text{SiNc(OR)}_2$ . These correspond to reversible one electron waves for  $\text{SiNc(OR)}_2$  and  $\text{SiPc(OR)}_2$ , but are two-electron waves for  $\text{RO(SiPcO)}_2\text{R}$ . The values of diffusion constants ( $D$ ,  $\text{cm}^2/\text{sec}$ ) computed from  $i(t)$  vs  $t^{-1/2}$  behavior (potential step experiment) for the neutral  $\text{SiPc(OR)}_2$ ,  $\text{RO(SiPcO)}_2\text{R}$ , and  $\text{SiNc(OR)}_2$  are  $5.7 \pm 1.1 \times 10^{-6} \text{ cm}^2/\text{s}$ ,  $2.2 \pm 0.3 \times 10^{-6} \text{ cm}^2/\text{s}$ , and  $5.2 \pm 0.4 \times 10^{-6} \text{ cm}^2/\text{s}$ , respectively. The compounds show absorptions corresponding to N bands, Soret bands, and intense absorption around 650-700 nm for  $\text{SiPc(OR)}_2$ , 600-650 nm for  $\text{RO(SiPcO)}_2\text{R}$ , and 750 - 800 nm for  $\text{SiNc(OR)}_2$  corresponding to Q bands. The corrected ecl maxima are around 684 nm for  $\text{SiPc(OR)}_2$  and 792 nm for  $\text{SiNc(OR)}_2$ . The ecl emission for  $\text{SiNc(OR)}_2$  is at the longest wavelength reported so far. No ecl was observed from the dimeric phthalocyanine.

#### ACKNOWLEDGMENT

We gratefully acknowledge partial support of this work by the Office of Naval Research (ONR), the Army Research Office, and by the Graduate School Annual Alumni Fund of Case Western Reserve University through a fellowship.

## REFERENCES

- (1) Lever, A. B. P. Adv. Inorg. Chem. Radiochem. 1965, 7, 27.
- (2) Grammatica, S.; Mort, J. Appl. Phys. Lett. 1981, 38, 445.
- (3) Mizuguchi, J. Jpn. J. Appl. Phys. 1981, 20, 293.
- (4) Kuznesof, P. M.; Nohr, R. S.; Wynne, K. J.; Kenney, M. E. J. Macromol. Sci.-Chem. 1981, A16, 299.
- (5) Lever, A. B. P.; Licoccia, S.; Ramaswamy, B. S.; Kandil, A.; Stynes, D. V. Inorg. Chim. Acta 1980, 51, 169.
- (6) (a) Van Ewyk, R. L.; Chadwick, A. V.; Wright, J.D. J. Chem. Soc. Faraday Trans. 1981, 77, 73; (b) Diel, B. N.; Inabe, T.; Lyding, J. W.; Schoch, Jr., K. F.; Kannewurf, C. R.; Marks, T. J. J. Am. Chem. Soc. 1983, 105, 1539.
- (7) Nohr, R. S.; Kuznesof, P. M.; Wynne, K. J.; Kenney, M. E.; Siebenman, P. G. J. Am. Chem. Soc. 1981, 103, 4371.
- (8) (a) Leempoel, P.; Castro-Acuna, M.; Fan, F.-R. F.; Bard, A. J. J. Phys. Chem., 1982, 86, 1396; (b) Jaeger, C. D.; Fan, F.-R. F.; Bard, A. J. J. Am. Chem. Soc. 1980, 102, 2592.
- (9) Shepard, Jr., V. R.; Armstrong, N. R. J. Phys. Chem. 1979, 83, 1268.
- (10) Krueger, P. C.; Kenney, M. E. J. Org. Chem. 1963, 28, 3379.
- (11) (a) Joyner, R. D.; Kenney, M. E. Inorg. Chem. 1962, 1, 717; (b) Hanack, M.; Mitulla, K.; Pawlowski, G.; Subramanian, L. R. J. Organometal. Chem. 1981, 204, 315.
- (12) Kovshev, E. I.; Puchnova, V. A.; Luk'yanets, E. A. J. Org. Chem. USSR (Eng. Trans.), 1971, 7, 364; Zh. Org. Khim. 1971, 7, 369.
- (13) Lowery, M. K.; Starshak, A. J.; Esposito, J. N.; Krueger, P. C.; Kenney, M. E. Inorg. Chem. 1965, 4, 128.
- (14) Douglass, S. L. Ph. D. Dissertation, Case Western Reserve University,



Cleveland, Ohio, 1975.

- (15) Janson, T. R. Ph. D. Dissertation, Case Western Reserve University, Cleveland, Ohio, 1971.
- (16) Mezza, T. M.; Armstrong, N. R.; Ritter, G. W., II; Iafalice, J. P.; Kenney, M. E. J. Electroanal. Chem. 1982, 137, 227.
- (17) Keszthelyi, C. P.; Tachikawa, H.; Bard, A. J. J. Am. Chem. Soc. 1972, 94, 1522.
- (18) Joyner, R. D.; Kenney, M. E. Inorg. Chem. 1962, 1, 236.
- (19) Pommier, J. C. Rev. Silicon, Germanium, Tin, Lead Compd. 1979, 4, 91.
- (20) Jansen, T. R.; Kane, A. R.; Sullivan, J. F.; Knox, K.; Kenney, M. E.; J. Am. Chem. Soc. 1969, 91, 5210.
- (21) (a) Clack, D. W.; Hush, N. S.; Woolsey, I. S. Inorg. Chim. Acta 1976, 19, 129; (b) Lever, A. B. P.; Wilshire, J. P.; Inorg. Chem. 1978, 17, 1145; (c) Lever, A. B. P.; Licoccia, S.; Magnell, K.; Minor, P. C.; Ramaswamy, B. S. Adv. in Chem. Series 1982, 201, 237.
- (22) Bard, A. J.; Faulkner, L. R. "Electrochemical Methods. Fundamentals and Applications," John Wiley and Sons, NY, 1980.
- (23) (a) Flanagan, J. B.; Margel, S.; Bard, A. J.; Anson, F. C. J. Am. Chem. Soc. 1978, 100, 4248; (b) Ammar, F.; Savéant, J. M.; J. Electroanal. Chem. 1973, 47, 215.
- (24) Mikhalenko, S. A.; Luk'yanets, E. A. J. Gen. Chem. USSR (Engl. Trans.) 1969, 39, 2495; Zh. Obshch. Khim. 1969, 39, 2554.
- (25) Vogler, A.; Kunkely, H. Inorg. Chim. Acta 1980, 44, L209.
- (26) Gouterman, M. "The Porphyrins," Dolphin, D. (ed), Academic Press, NY 1978, Vol. III, Ch. 1.
- (27) (a) Kane, A. R.; Sullivan, J. F.; Kenny, D. H.; Kenney, M. E. Inorg. Chem. 1970, 9, 1445; (b) Hush, N. S.; Woolsey, I. S.; Mol. Phys. 1971, 21, 465.

- (28) Kasha, M. Radiation Res. 1963, 20, 55.
- (29) Tokel, N. E.; Keszthelyi, C. P.; Bard, A. J. J. Am. Chem. Soc. 1972, 94, 4872.
- (30) Faulkner, L. R.; Tachikawa, H.; Bard, A. J. J. Am. Chem. Soc. 1972, 94, 691.
- (31) (a) Tokel, N. E.; Bard, A. J. J. Am. Chem. Soc. 1972, 94, 2862; (b) Chang, J.; Hercules, D. M.; Roe, D. K. Electrochim. Acta 1968, 13, 1197.
- (32) Stadelmann, H. R.; J. Luminescence 1972, 5, 171.
- (33) Doblhofer, K.; Gerischer, H. Electrochim. Acta 1975, 20, 215.

# FIGURE CAPTIONS

- Figure 1 The structures of  $\text{SiPc}[\text{OSi}(\text{n-C}_6\text{H}_{13})_3]_2$  and  $\text{SiNc}[\text{OSi}(\text{n-C}_6\text{H}_{13})_3]_2$ .
- Figure 2 The infrared spectrum of  $\text{SiNc}(\text{OH})_2$  in a Nujol mull (KBr plates).
- Figure 3 The 200 MHz NMR spectra of (a)  $\text{SiPc}[\text{OSi}(\text{n-C}_6\text{H}_{13})_3]_2$  and (b)  $\text{SiNc}[\text{OSi}(\text{n-C}_6\text{H}_{13})_3]_2$  in  $\text{CDCl}_3$ .
- Figure 4 Cyclic voltammogram of (a) 2.2 mM  $\text{SiPc}[\text{OSi}(\text{n-C}_6\text{H}_{13})_3]_2$  and (b) 1.4 mM  $\text{SiNc}[\text{OSi}(\text{n-C}_6\text{H}_{13})_3]_2$ , and (c) 1.0 mM  $(\text{n-C}_6\text{H}_{13})_3\text{SiO}(\text{SiPcO})_2\text{Si}(\text{n-C}_6\text{H}_{13})_3$  in  $\text{CH}_2\text{Cl}_2$ , 0.1 M TBAP. Scan rate = 20 mV/s.
- Figure 5 (a) ac voltammogram of  $\text{SiPc}[\text{OSi}(\text{n-C}_6\text{H}_{13})_3]_2$ , 2.2 mM, in  $\text{CH}_2\text{Cl}_2$ , 0.1 M TBAP; (b) ac voltammogram of  $\text{SiNc}[\text{OSi}(\text{n-C}_6\text{H}_{13})_3]_2$ , 1.4 mM, in  $\text{CH}_2\text{Cl}_2$ , 0.1 M TBAP; (c) ac voltammogram of  $(\text{n-C}_6\text{H}_{13})_3\text{SiO}(\text{SiPcO})_2\text{Si}(\text{n-C}_6\text{H}_{13})_3$ , 1.0 mM, in  $\text{CH}_2\text{Cl}_2$ , 0.1 M TBAP. For all: ac frequency = 50 Hz; scan rate = 2 mV/s; in phase (—); phase angle (----).
- Figure 6 (a) Second harmonic ac voltammogram for the solution in Figure 5a; (b) same for solution in Figure 5b; (c) same for solution Figure 5c. For all: ac fundamental frequency = 200 Hz; scan rate = 2 mV/s.
- Figure 7 The electronic absorption spectra of  $1.1 \times 10^{-5}$  M  $\text{SiPc}[\text{OSi}(\text{n-C}_6\text{H}_{13})_3]_2$  (----),  $1.2 \times 10^{-5}$  M  $(\text{n-C}_6\text{H}_{13})_3\text{SiO}(\text{SiPcO})_2\text{Si}(\text{n-C}_6\text{H}_{13})_3$  (—) and  $2.5 \times 10^{-6}$  M  $\text{SiNc}[\text{OSi}(\text{n-C}_6\text{H}_{13})_3]_2$  (—) in  $\text{CH}_2\text{Cl}_2$ .
- Figure 8 (a) Fluorescence spectrum (uncorrected) of  $1.1 \times 10^{-5}$  M  $\text{SiPc}[\text{OSi}(\text{n-C}_6\text{H}_{13})_3]_2$  in  $\text{CH}_2\text{Cl}_2$ . For excitation spectrum (—), the emission was monitored at 684 nm. For the emission spectrum (----), the excitation was at 350 nm. Bandwidth = 5 nm; (b) fluorescence spectrum (uncorrected) for  $1.0 \times 10^{-5}$  M  $\text{SiNc}[\text{OSi}(\text{n-C}_6\text{H}_{13})_3]_2$  in  $\text{CH}_2\text{Cl}_2$ . For excitation spectrum (—), emission was monitored at 792 nm. For emission spectrum (----), excitation was at 356 nm. Bandwidth = 5 nm.

9 Ecl spectra of (a)  $\text{SiPc}[\text{OSi}(\text{n-C}_6\text{H}_{13})_3]_2$ ,  $\lambda_{\text{max}} = 725 \text{ nm}$ , and (b)  $\text{SiNc}[\text{OSi}(\text{n-C}_6\text{H}_{13})_3]_2$ ,  $\lambda_{\text{max}} = 828 \text{ nm}$ . Concentrations are 2 mM and 1.4 mM, respectively, in  $\text{CH}_2\text{Cl}_2$ , 0.1 M TBAP. Pt electrode pulsed from +1.0 V to -1.2 V, and +0.6 V to -1.3 V vs. AgRE, respectively, at 0.5 Hz. The greater emission for both occurred on the cathodic pulse.

Table 1



Values of  $E^\circ$  (V vs. SCE) for  $\text{SiPc(OR)}_2$ ,  $\text{RO(SiPcO)}_2\text{R}$ , and  $\text{SiNc(OR)}_2$  in  $\text{CH}_2\text{Cl}_2$

	$\text{SiPc(OR)}_2$ (V vs. SCE)	$\text{SiNc(OR)}_2$ (V vs. SCE)	$\text{RO(SiPcO)}_2\text{R}$ (V vs. SCE)
2nd ox.	--	+1.24	+1.20
1st ox.	+1.00	+0.58	+0.71
1st red.	-0.90	-1.01	-0.81
2nd red.	-1.48	-1.55	-1.21
$-\Delta H^\circ$	$1.8 \pm 0.1$ eV	$1.5 \pm 0.1$ eV	--
singlet emission energy	1.86 eV	1.61 eV	--

Table 2

Q-Band Absorption and Emission Maxima (nm) of SiPc(OR)<sub>2</sub> and SiNc(OR)<sub>2</sub>

	SiPc(OR) <sub>2</sub>		SiNc(OR) <sub>2</sub>	
	Absorption	Emission	Absorption	Emission
$S_0(0) \leftrightarrow S_1(0)$	668	668	772	772
$S_0(0) \leftrightarrow S_1(1)$	638	708	732	790
$S_0(0) \leftrightarrow S_1(2)$	612	741	704	--
$S_0(0) \leftrightarrow S'_1(0)$	601	--	686	--
$S_0(0) \leftrightarrow S'_1(1)$	575	--	655	--
$S_0(0) \leftrightarrow S'_1(2)$	557	--	636	--






Fig. 2







Fig. 3

Fig. 4






Fig. 5




Fig. 6






Fig. 7

(a)



(b)







Fig. 9

DL/413/83/01  
GEN/413-2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 413 800 N. Quincy Street Arlington, Virginia 22217	2	Naval Ocean Systems Center Attn: Technical Library San Diego, California 92152	1
ONR Pasadena Detachment Attn: Dr. R. J. Marcus 1030 East Green Street Pasadena, California 91106	1	Naval Weapons Center Attn: Dr. A. B. Amster Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1	Dean William Tolles Naval Postgraduate School Monterey, California 93940	1
Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Mr. Vincent Schaper DTNSRDC Code 2830 Annapolis, Maryland 21402	1
DTNSRDC Attn: Dr. G. Bosmajian Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1	Mr. A. M. Anzalone Administrative Librarian PLASTEC/ARRADCOM Bldg 3401 Dover, New Jersey 07801	1



TECHNICAL REPORT DISTRIBUTION LIST, 356B

Dr. C. L. Schilling  
Union Carbide Corporation  
Chemical and Plastics  
Tarrytown Technical Center  
Tarrytown, New York

Dr. A. G. MacDiarmid  
Department of Chemistry  
University of Pennsylvania  
Philadelphia, Pennsylvania 19174

Dr. E. Fischer, Code 2853  
Naval Ship Research and  
Development Center  
Annapolis, Maryland 21402

Dr. H. Allcock  
Department of Chemistry  
Pennsylvania State University  
University Park, Pennsylvania 16802

Dr. M. Kenney  
Department of Chemistry  
Case Western University  
Cleveland, Ohio 44106

Dr. R. Lenz  
Department of Chemistry  
University of Massachusetts  
Amherst, Massachusetts 01002

Dr. M. David Curtis  
Department of Chemistry  
University of Michigan  
Ann Arbor, Michigan 48105

NASA-Lewis Research Center  
Attn: Dr. T. T. Serafini, MS 49-1  
21000 Brookpark Road  
Cleveland, Ohio 44135

Dr. J. Griffith  
Naval Research Laboratory  
Chemistry Section, Code 6120  
Washington, D.C. 20375

Professor G. Wnek  
Department of Materials Science  
and Engineering  
Massachusetts Institute of Technology  
Cambridge, Massachusetts 02139

Dr. R. Soulen  
Contract Research Department  
Pennwalt Corporation  
900 First Avenue  
King of Prussia, Pennsylvania 19406

Dr. G. Goodman  
Globe-Union Incorporated  
5757 North Green Bay Avenue  
Milwaukee, Wisconsin 53201

Dr. Martin H. Kaufman  
Code 38506  
Naval Weapons Center  
China Lake, California 93555

Dr. C. Allen  
Department of Chemistry  
University of Vermont  
Burlington, Vermont 05401

Professor R. Drago  
Department of Chemistry  
University of Florida  
Gainesville, Florida 32611

Dr. D. L. Venezky  
Code 6130  
Naval Research Laboratory  
Washington, D.C. 20375

Professor T. Katz  
Department of Chemistry  
Columbia University  
New York, New York 10027

Professor James Chien  
Department of Chemistry  
University of Massachusetts  
Amherst, Massachusetts 01002

Professor J. Salamone  
Department of Chemistry  
University of Lowell  
Lowell, Massachusetts 01854

Dr. S. Cooper  
Department of Chemistry  
University of Wisconsin  
750 University Avenue  
Madison, Wisconsin 53706

DL/413/83/01  
356B/413-2

TECHNICAL REPORT DISTRIBUTION LIST, 356B

Professor D. Grubb  
Department of Materials Science  
and Engineering  
Cornell University  
Ithaca, New York 14853

Professor T. Marks  
Department of Chemistry  
Northwestern University  
Evanston, Illinois 60201

Professor C. Chung  
Department of Materials Engineering  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Professor Malcolm B. Polk  
Department of Chemistry  
Atlanta University  
Atlanta, Georgia 30314

Dr. D. B. Cotts  
SRI International  
333 Ravenswood Avenue  
Menlo Park, California 94205

Dr. Kurt Baum  
Fluorochem, Inc.  
680 S. Ayon Avenue  
Azusa, California 91702

Professor H. Hall  
Department of Chemistry  
University of Arizona  
Tucson, Arizona 85721

Professor G. Whitesides  
Department of Chemistry  
Harvard University  
Cambridge, Massachusetts 02138

Professor H. Ishida  
Department of Macromolecular Science  
Case Western University  
Cleveland, Ohio 44106

Dr. K. Paciorek  
Ultrasystems, Inc.  
P.O. Box 19605  
Irvine, California 92715

Professor D. Seyferth  
Department of Chemistry  
Massachusetts Institute of Technology  
Cambridge, Massachusetts 02139

Dr. G. Bryan Street  
IBM Research Laboratory, K32/281  
San Jose, California 95193

**DAT  
FILM**